


Carol A. See

Name of Person Signing



Signature

**PATENT**  
**Docket No. GC626-2**

**Patent Application for**  
**Mass Spectrometric Analysis Of Biopolymers**

Inventors: David A. Estell, Grant C. Ganshaw, Christian Paech, and  
Sigrid Paech

**CROSS-REFERENCE TO RELATED APPLICATIONS**

Pursuant to 35 U.S.C. §119(e), the present application claims benefit of and  
priority to USSN 60/228,198, entitled "Mass Spectrometric Analysis of Biopolymers,"  
filed August 25, 2000, by Christian Paech et al.

**FIELD OF THE INVENTION**

The present invention relates to the analysis of biopolymers in crude  
solutions. In particular, the invention relates to the determination, quantitation, and  
identification of biopolymers, such as polypeptides and oligonucleotides, using mass  
spectroscopic data obtained from fractionated mixtures.

**REFERENCES**

Allen G (1989) *Sequencing of Proteins and Peptides*. 2nd edn. Elsevier,  
Amsterdam.

Bairoch A, Apweiler R (2000) The SWISS-PROT protein sequence database  
and its supplement TrEMBL in 2000. *Nucleic Acids Res* **28**:45-48.

Burks C, et al. (1990) GenBank: current status and future directions.  
*Methods Enzymol* **183**:3-22.

Chowdhury SK et al. (1995) Examination of Recombinant Truncated Mature  
Human Fibroblast Collagenase by Mass Spectrometry: Identification of Differences  
with the Published Sequence and Determination of Stable Isotope Incorporation.  
*Rapid Communications in Mass Spectrometry* **9**:563-569.



Christianson T, Paech C (1994) Peptide mapping of subtilisins as a practical tool for locating protein sequence errors during extensive protein engineering projects. *Anal Biochem* **223**:119-129.

5 Corthals G.L., et al. (1999) Identification of proteins by mass spectrometry, in *Proteome research: 2D gel electrophoresis and detection methods*, Ed. Rabilloud, T., Springer, New York, pp. 197-231.

Deutscher MP, ed (1990) *Guide to Protein Purification*. Academic Press, New York.

10 George DG, et al. (1996) PIR-International Protein Sequence Database. *Methods Enzymol* **266**:41-59.

Goddette DW, et al. (1992) The crystal structure of the *Bacillus lentus* alkaline protease, subtilisin BL, at 1.4 Å resolution. *J Mol Biol* **228**:580-595.

15 Guermant C, et al. (2000) Under proper control, oxidation of proteins with known chemical structure provides an accurate and absolute method for the determination of their molar concentration. *Anal Biochem* **277**:46-57.

Gygi SP, et al. (1999) Quantitative analysis of complex protein mixtures using isotope-coded affinity tags. *Nat Biotechnol* **17**:994-999.

Hancock WS, ed (1996) *New Methods in Peptide Mapping for the Characterization of Proteins*. CRC Press, Boca Raton.

20 Hsia C, et al. (1996) Active-site titration of serine proteases using a fluoride ion selective electrode and sulfonyl fluoride inhibitors. *Anal Biochem* **242**:221-227.

Janson JC, Rydén L, eds (1998) *Protein Purification*. 2nd edn. Wiley-Liss, New York.

25 Kahn P, Cameron G (1990) EMBL Data Library. *Methods Enzymol* **183**:23-31.

Kellner R, Lottspeich F, Meyer HE, eds (1999) *Microcharacterization of Proteins*. 2nd edn. Wiley-VCH, Weinheim.

Kunst F, et al. (1997) The complete genome sequence of the gram-positive bacterium *Bacillus subtilis*. *Nature* **390**:249-256.

30 Lahm HW, Langen H (2000) Mass spectrometry: a tool for the identification of proteins separated by gels. *Electrophoresis* **21**:2105-2114.

Matsudaira P, ed (1993) *A Practical Guide to Protein and Peptide Purification for Microsequencing*. 2nd edn. Academic Press, San Diego.

35 Oda Y, et al. (1999) Accurate quantitation of protein expression and site-specific phosphorylation. *Proc Natl Acad Sci USA* **96**:6591-6596.

Pace CN, et al. (1995) How to measure and predict the molar absorption coefficient of a protein. *Protein Sci* 4:2411-2423.

Scopes R (1994) *Protein Purification*. 3rd edn. Springer-Verlag, New York.

Stocklin et al., (1997) A Stable Isotope Dilution Assay for the In Vivo

5 Determination of Insulin Levels in Humans by Mass Spectrometry. *Diabetes* 46:44-50.

## BACKGROUND OF THE INVENTION

Protein concentration determination is at the heart of any study concerned  
10 with the catalytic efficiency of an enzyme. Even for highly purified enzymes the choice of first-principle methods for accurately measuring molar concentrations is restricted to a few techniques (amino acid, total nitrogen, and absorbance measurement (Pace et al., 1995), titration of oxidized sulfur (Guermant et al., 2000). For enzymes in crude solution the options are even smaller and techniques are  
15 much more elaborate (e.g., active-site titrations involving the stoichiometric release of a reporter group, enzyme-linked immunosorbent assay (ELISA), densitometry after sodium dodecylsulfate polyacrylamide gel electrophoresis (SDS-PAGE)). Catalytic rate assays while highly specific for an enzyme and often quantitative in nature presuppose validation with purified enzyme which in turn requires first-principle  
20 methods for accurate mass quantitation.

The determination of the concentration of a specific protein among other proteins in crude solution, such as a fermenter broth, is a formidable challenge. Even more demanding is the task of verifying the presence of a specific protein and the quantitation of this protein in a cell or tissue extract without knowing the  
25 properties of the protein and ever having seen it before.

Most methods for estimating protein concentration are built on general properties of proteins, e.g., the chemistry and light absorbance of aromatic side chains and the peptide bond, and the binding affinity for chromophores. More specific techniques, e.g. immunoassay and active site titration, require some prior  
30 knowledge of the targeted protein. All such methods, however, suffer from interferences, as the extensive literature on protein assays documents, and none of the methods takes advantage of that one unique feature that differentiates non-identical proteins, the amino acid sequence. On that level there is no interference possible.

35 The use of isotopically labeled biopolymers to investigate cellular processes is not new. For example, Chowdhury et al. used mass spectrometry and isotopically

labeled analogs to investigate the molecular weight of truncated mature collagenase, and Stocklin et al. have investigated human insulin concentration in serum samples that had been extracted and purified. Neither one discuss the use of crude solutions to determine biopolymer concentration without prior isolation of the biopolymer.

5           The present invention makes use of the subunit sequence as a unique tag of a biopolymer (e.g., the amino acid sequence of a specific protein), that can be exploited for determining the concentration in crude solutions.

## SUMMARY OF THE INVENTION

10           The present invention addresses the need for a straightforward and rapid technique for determining the specific concentration of one or more biopolymers (e.g., proteins, oligonucleotides, etc.) in a mixture, e.g., a cell-free culture fluid, a cell extract, or the entire complement of proteins in a cell or tissue.

15           The present invention additionally provides a method for identifying a biopolymer fragment (e.g., peptide, oligonucleotide, etc.) derived from a larger biopolymer added to a solution that otherwise lacks such a biopolymer or fragment.

In one of its aspects, the present invention provides a method for determining the absolute quantity of a target polypeptide, such as a selected protein, in a crude solution or mixture, comprising the steps of:

- 20           (a) adding a known quantity of an analog of the target polypeptide to the solution or mixture;
- (b) treating the target polypeptide and analog in the solution or mixture with a fragmenting activity (e.g., a protease) to generate a plurality of corresponding peptide pairs;
- 25           (c) resolving the peptide content of the solution or mixture;
- (d) determining by mass spectrometric analysis the ratio of a selected target peptide to its corresponding analog peptide; and
- (e) calculating, from the ratio and the known quantity of the analog, the quantity of the target polypeptide in the solution or mixture.

30           The solution or mixture can be, for example, a crude fermenter solution, a cell-free culture fluid, a cell extract, or a mixture comprising the entire complement of proteins in a cell or tissue.

35           Another aspect of the present invention provides a method for determining the absolute quantity of a target polynucleotide in a crude solution, comprising the steps of:

(a) adding a known quantity of an analog of the target polynucleotide to the solution;

(b) treating the target polynucleotide and analog with a fragmenting activity (e.g., a restriction enzyme) to generate a plurality of corresponding polynucleotide-fragment pairs;

(c) resolving the polynucleotide-fragment content of the mixture;

(d) determining by mass spectrometric analysis the ratio of a selected target polynucleotide fragment to its corresponding analog fragment; and

(e) calculating, from the ratio and the known quantity of the analog, the quantity of the target oligonucleotide in the mixture.

In one embodiment, the target polynucleotide is an oligonucleotide.

Yet a further aspect of the present invention provides a method for verifying the presence and, optionally, determining the absolute quantity of a selected putative polypeptide, such as a protein, in a mixture containing a plurality of isotope-labeled cellular proteins from a selected cell type. One embodiment of the method includes the steps of:

selecting a putative polypeptide potentially present in said mixture;

generating a theoretical fragmentation of the putative polypeptide;

selecting a theoretical fragment from the theoretical fragmentation;

producing a peptide having an amino acid sequence corresponding to the theoretical fragment;

adding a known amount of the produced peptide as an internal standard to the mixture;

treating the mixture with a proteolytic activity;

resolving the cellular polypeptide fragments along with the internal standard and analyzing the same by mass spectrometry to provide a mass spectrograph;

locating a peak pair from the mass spectrograph comprised of a peak representing the internal standard and a peak representing a cellular polypeptide fragment corresponding to the internal standard, thereby verifying the presence of the putative polypeptide;

optionally, upon verifying the presence of the putative polypeptide, determining the ratio of internal standard to its corresponding cellular polypeptide fragment; and,

calculating, from the ratio and the known quantity of the internal standard, the absolute quantity of the putative polypeptide in the mixture.

The putative polypeptide can be derived, for example, from a database of sequence information.

Preferably, in connection with the fragmentation step, the fragmentation of the cellular polypeptide is determined to be substantially complete with respect to the cellular polypeptide fragment corresponding to the internal standard.

One embodiment provides the additional steps of:

after determining the absolute quantity of the putative polypeptide in the mixture, growing the selected cell type under a set of defined conditions, querying an extract from the grown cell type for the presence, for an increase or decrease of the absolute concentration of the putative polypeptide by mixing the extract with a known amount of the isotope-labeled mixture as a new internal standard;

treating the extract with a proteolytic activity;

resolving the polypeptide fragment content of the extract and analyzing the same by mass spectrometry to provide a mass spectrograph;

locating a peak pair from said mass spectrograph comprised of a peak representing the new internal standard and a peak representing a cellular polypeptide fragment corresponding to the new internal standard, thereby verifying the presence of the putative polypeptide;

optionally, upon verifying the presence of the putative polypeptide, determining the ratio of the new internal standard to its corresponding cellular polypeptide fragment; and,

calculating, from the ratio and the known quantity of the internal standard, the absolute quantity of the putative polypeptide in the extract.

In another of its aspects, the present invention provides a cell-culture extract, derived from a selected microorganism grown on media enriched in a specific isotope, said extract containing a known amount of a metabolically labeled polypeptide determined by a peptide-separation technique in combination with mass spectroscopy.

A further aspect of the present invention provides a method for determining the identity of a target polypeptide fragment in a solution, comprising the steps of:

(a) adding an analog of the target polypeptide and the target polypeptide to the solution, in a selected fixed analog:target ratio;

(b) treating the target polypeptide and analog with a fragmenting activity to generate a plurality of corresponding peptide pairs;

- (c) resolving the peptide content of the solution;
- (d) identifying by mass spectrometric analysis those fragment pairs that exhibit the selected ratio; and, optionally,
- (e) determining the amino acid sequence of the fragment pairs identified in step (d).

In one embodiment, the target polypeptide is a protein.

In another embodiment, the crude solution contains a plurality of different proteins. For example, the solution can be a crude fermenter solution, a cell-free culture fluid, a cell extract, a mixture comprising the entire complement of proteins in a cell or tissue, etc.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the scope and spirit of the invention will become apparent to one skilled in the art from this detailed description.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. UV traces of a tryptic co-digest of  $^{15}\text{N}$ -subtilisin-DAI, indexed ( $^{15}\text{N}$ ), and subtilisin, indexed (s). Peptide numbering refers to Table I.

Figure 2. Total ion current chromatogram of selected peptides in Figure 1. (A) Peptide 3 of subtilisin (3 (s), upper panel) and peptide 3 of  $^{15}\text{N}$ -subtilisin-DAI (3 ( $^{15}\text{N}$ ), lower panel). (B) TIC of peptides 5, 6, and 9 of the co-digest of  $^{15}\text{N}$ -subtilisin-DAI, indexed ( $^{15}\text{N}$ ), and subtilisin, indexed (s). Sequence differences between subtilisin-DAI and subtilisin reside on peptide 5 (N74D) and 6 (S101A, V102I). Amino acid sequence numbering is linear.

Figure 3. Rapid tryptic digest of subtilin-DAI and  $^{15}\text{N}$ -subtilisin-DAI and separation of peptides by RP-HPLC on a 2.0x50 mm C18 column (Jupiter, by Phenomenex). The quantitation by TIC peak area integration of corresponding peaks gave the result expected from enzyme activity assays and active site titrations (see Figures 1 and 2).

Figure 4. (A) SDS-PAGE of a fermentation broth concentrate of unknown origin. (B) This material spiked with a known amount of  $^{15}\text{N}$ -labeled purified subtilisin BPN'-

Y217L and was digested with trypsin. The peptide mixture was separated by RP-HPLC on a C18 column (2.1 x 150 mm) and the eluate was recorded at 215 nm.

Sub A1  
Sub B1

Figure 5. Total ion current chromatogram of peptides 1, 2, and 3 from Figure 3. (1) Mass 980.6 (1+), left trace; mass 991.5 (1+), right trace, corresponding to tryptic peptide SSLENTTTK of BPN' and containing 11 nitrogen atoms. (2) Mass 765.6(2+), left trace; mass 775.6 (2+), right trace corresponding to tryptic peptide APALHSQGYTGSNVK of BPN' and containing 20 nitrogen atoms. 'x' is an unrelated peptide. (3) Mass 627.0 (2+), left trace; mass 636.4(2+), right trace corresponding to tryptic peptide HPNWTNTQVR of BPN' and containing 19 nitrogen atoms.

Figure 6. Table I.: Sequence comparison, m/z values, and ratios of integrated TIC peak areas and UV absorbance peak areas for chromatogram in Figure 1. The concentration measured by the co-digest technique for subtilisin and subtilisin-DAI was 8.15 and 7.13 mg/ml, respectively, while the given concentration (established by independent methods) was 7.99 and 7.03mg/ml, respectively.

Figure 7. Table II. Determination of concentration, activity and conversion factor for subtilisin-DAI variants determined by peptide mapping (<sup>15</sup>N-isotope method) and by active site titration with a calibrated mung bean inhibitor solution using as internal standard a previously calibrated solution of subtilisin-DAI (Hsia et al., 1996). The range of target protein concentrations was 2 to 5 µg · ml<sup>-1</sup>.

## DETAILED DESCRIPTION OF THE INVENTION

The invention will now be described in detail by way of reference only using the following definitions and examples. All patents and publications, including all sequences disclosed within such patents and publications, referred to herein are expressly incorporated by reference.

The present invention provides methods for the quantitation of biopolymers in crude, i.e., unpurified, solutions.

### Definitions

Unless defined otherwise herein, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Singleton, *et al.*, DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY, 2d Ed., John Wiley and Sons, New York (1994), and Hale & Marham, THE HARPER COLLINS DICTIONARY OF BIOLOGY, Harper Perennial, NY (1991)





provide one of skill with a general dictionary of many of the terms used in this invention. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described. Numeric ranges are inclusive of the numbers defining the range. Unless otherwise indicated, nucleic acids are written left to right in 5' to 3' orientation; amino acid sequences are written left to right in amino to carboxy orientation, respectively. The headings provided herein are not limitations of the various aspects or embodiments of the invention which can be had by reference to the specification as a whole. Accordingly, the terms defined immediately below are more fully defined by reference to the specification as a whole.

Biopolymer

The term "biopolymer" as used herein means any large polymeric molecule produced by a living organism. Thus, it refers to nucleic acids, polynucleotides, polypeptides, proteins, polysaccharides, carbohydrates, lipids and analogues thereof. The terms "biopolymer" and "biomolecule" are used interchangeably herein.

Isolated

As used herein an "isolated" biomolecule (such as a nucleic acid or protein) has been substantially separated or purified away from other biological components in the cell of the organism in which the component naturally occurs, i.e., other chromosomal and extrachromosomal DNA and RNA, and proteins. Nucleic acids and proteins which have been "isolated" thus include nucleic acids and proteins purified by standard purification methods. The term also embraces nucleic acids and proteins prepared by recombinant expression in a host cell as well as chemically synthesized nucleic acids.

Polypeptide or Protein

A macromolecule composed of one to several polypeptides. Each polypeptide consists of a chain of amino acids linked together by covalent (peptide) bonds. They are naturally-occurring complex organic substances composed essentially of carbon, hydrogen, oxygen and nitrogen, plus sulphur or phosphorus, which are so associated as to form sub-microscopic chains, spirals or plates and to which are attached other atoms and groups of atoms in a variety of ways. A protein may comprise one or multiple polypeptides linked together by disulfid bonds. Examples of the protein include, but are not limited to, antibodies, antigens, ligands,

receptors, etc. The terms "polypeptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues.

As the description of this invention proceeds, it will be seen that mixtures are produced which may contain individual components containing 100 or more amino acid residues or as few as one or two such residues. Conventionally, such low molecular weight products would be referred to as amino acids, dipeptides, tripeptides, etc. However, for convenience herein, all such products will be referred to as polypeptides since the mixtures which are prepared for mass spectrometric analysis contain such components together with products of sufficiently high molecular weight to be conventionally identified as polypeptides.

Polypeptides may contain amino acids other than the 20 gene encoded amino acids. "Polypeptide(s)" include those modified either by natural processes, such as processing and other post-translational modifications, but also by chemical modification techniques. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known to those of skill in the art. Polypeptides may be branched or cyclic, with or without branching. Cyclic, branched and branched circular polypeptides may result from post-translational natural processes and may be made by entirely synthetic methods, as well.

Peptide or oligopeptide

A linear molecule composed of two or more amino acids linked by covalent (peptide) bonds. They are called dipeptides, tripeptides and so forth, according to the number of amino acids present. These terms may be used interchangeably with polypeptide. See above.

Polynucleotide

A chain of nucleotides in which each nucleotide is linked by a single phosphodiester bond to the next nucleotide in the chain. They can be double- or single-stranded. The term is used to describe DNA or RNA.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotide(s)" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA

that may be single-stranded or, more typically, double-stranded, or a mixture of single- and double-stranded regions. The RNA may be a mRNA.

As used herein, the term "polynucleotide(s)" also includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotide(s)" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as 4-acetylcytosine, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term "polynucleotide(s)" as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including, for example, simple and complex cells.

The length of the polynucleotides may be 10 kb. In accordance with one embodiment of the present invention, the length of a polynucleotide is in the range of about 50 bp to 10 Kb, preferably, 100 bp to 1.5 kb.

#### Oligonucleotide

A short molecule (usually 6 to 100 nucleotides) of single-stranded DNA. "Oligonucleotide(s)" refer to short polynucleotides, i.e., less than about 50 nucleotides in length. In a preferred embodiment, the oligonucleotides can be of any suitable size, and are preferably 24-48 nucleotides in length. In accordance with another embodiment of the present invention, the length of a synthesized oligonucleotide is in the range of about 3 to 100 nucleotides. In accordance with a further embodiment of the present invention, the length of the oligonucleotide is in the range of about 15 to 20 nucleotides.

Size separation of the cleaved fragments is performed using 8 percent polyacrylamide gel described by Goeddel et al., Nucleic Acids Res., 8:4057 (1980).

#### Restriction enzyme

Restriction enzyme and restriction endonuclease are used interchangeably herein and refer to a protein that recognizes specific, short nucleotide sequences and cuts the DNA at those sites. There are three types of restriction endonuclease enzymes:

Type I: Cuts non-specifically a distance greater than 1000 bp from its recognition sequence and contains both restriction and methylation activities.



Type II: Cuts at or near a short, and often palindromic recognition sequence. A separate enzyme methylates the same recognition sequence. They may make the cuts in the two DNA strands exactly opposite one another and generate blunt ends, or they may make staggered cuts to generate sticky ends. The type II restriction enzymes are the ones commonly exploited in recombinant DNA technology.

Type III: Cuts 24-26 bp downstream from a short, asymmetrical recognition sequence. Requires ATP and contains both restriction and methylation activities.

The present invention contemplates the fragmentation of polynucleotides with restriction enzymes. In a preferred embodiment the restriction enzyme is a Type II. The fragment polynucleotides are then resolved into individual components based on size.

#### The Invention

In one of its aspects, the present invention makes use of the biomolecule (e.g., amino acid or nucleotide) sequence as a unique tag of a specific biopolymer (e.g., polypeptide or polynucleotide) that can be exploited for determining biopolymer concentration or identity in crude solutions, e.g., a crude fermenter solution, a cell-free culture fluid, a cell or tissue extract, etc. In one general embodiment, a target biomolecule is selected for analysis and an analog thereof is generated. The analog is purified and calibrated, and a known amount is added as an internal standard to the solution to be assayed. The biopolymers of the mixture are then fragmented, e.g., by proteolytic digestion for proteins, and the resulting biomolecule-fragments are resolved, e.g., by way of chromatography. One or more corresponding biomolecule-fragments pairs are then identified and analyzed by selected ion monitoring of a mass spectrometer.

According to one general embodiment, a target polypeptide is selected for analysis and an analog of the target polypeptide is generated. The target protein can be, for example, a protein that is known to be in a mixture, a putative protein (e.g., derived from a genome database search) that is potentially present in a mixture, or a known or putative protein segment or fragment (peptide). The analog of the target polypeptide can be the target polypeptide itself or a unique segment or fragment (peptide) of the target polypeptide. One or the other of the target polypeptide and analog is labeled so that the two can be distinguished from one another in subsequent mass analysis. The analog is purified and its absolute

quantity is determined in a solid quantity or in a solution by standard techniques (the analog is now said to be 'calibrated'), and a known amount is employed as an internal standard in the solution to be assayed. The polypeptides of the mixture are treated with a fragmenting activity, and the peptide components of the mixture are then resolved. Corresponding peptide pairs are then analyzed by selected ion monitoring of a mass spectrometer. Peak area integration of such peptide pairs provides a direct measure for the amount of target polypeptide in the crude solution.

According to another embodiment, a target polynucleotide is selected for analysis and an analog of the target polynucleotide is generated. The target polynucleotide can be, for example, a gene sequence that is known to be in a mixture, a putative gene (e.g., derived from a genome database search) that is potentially present in a mixture, or a known or putative polynucleotide or fragment (oligonucleotide). The analog of the target polynucleotide can be the target polynucleotide itself or a unique segment or fragment (oligonucleotide) of the target polynucleotide. One or the other of the target polynucleotide and analog is labeled so that the two can be distinguished from one another in subsequent mass analysis. The analog is purified and its absolute quantity is determined in a solid quantity or in a solution by standard techniques (the analog is now said to be 'calibrated'), and a known amount is employed as an internal standard in the solution to be assayed. The polynucleotides of the mixture are treated with a fragmenting activity, and the oligonucleotide components of the mixture are then resolved. Corresponding nucleotide-fragment pairs are then analyzed by selected ion monitoring of a mass spectrometer. Peak area integration of such nucleotide-fragment pairs provides a direct measure for the amount of target polynucleotide in the crude solution.

In yet another embodiment, the biomolecule analog is labeled with a suitable stable isotope and calibrated. The sample containing (or suspected of containing) the biomolecule of interest is aliquoted out such that the final concentration (after addition of the analog) in each aliquot is the same. Then decreasing amounts of the known labeled biomolecule analog is added to each aliquot. Each aliquot is subjected to mass spectrometry and their spectra analyzed for peaks corresponding to the labeled and unlabeled biomolecule of interest. Corresponding biomolecule peaks of the same magnitude, i.e., where the peak area ratio of labeled:unlabeled biomolecule equals one, indicates that the concentrations of each are the same. Thus, one is able to determine the concentration of the unlabeled biomolecule of interest from the sample with the known concentration of the labeled analog when the ratio equals one.

In a further embodiment, neither the biomolecule of interest nor the analog are labeled with a stable isotope. A known quantity of the analog is added in decreasing amounts to aliquots of the sample to be analyzed to yield a contaminated sample. The contaminated sample is treated with a fragmenting activity, and the biomolecule components of the mixture resolved. The resolved biomolecule-fragments, i.e., the corresponding biomolecule-fragment pairs, are then analyzed by mass spectrometry. The contribution of the unlabeled contaminant will decrease as its concentration in the sample of interest decreases. At some concentration the contribution of the unlabeled analog to the spectral analysis becomes negligible and the concentration of the biomolecule of interest can be determined. The concentration of the biomolecule of interest is determined by the intensity of the signal when the contribution of the analog is negligible and known concentration of the analog.

#### Isotope Labeling of Proteins

Labeling of the target or analog can be effected by any means known in the art. For example, a labeled protein or peptide can be synthesized using isotope-labeled amino acids or peptides as precursor molecules. Preferred labeling techniques utilize stable isotopes, such as  $^{18}\text{O}$ ,  $^{15}\text{N}$ ,  $^{13}\text{C}$ , or  $^2\text{H}$ , although others may be employed. Metabolic labeling can also be used to produce labeled proteins and peptides. For example, cells can be grown on a media containing isotope-labeled precursor molecules. Particularly, an organism can be grown on  $^{15}\text{N}$ -labeled organic or inorganic material, such as urea or ammonium chloride, as the sole nitrogen source. See Example 5.

In a preferred method, biopolymers are labeled with  $^{15}\text{N}$ . The following is a preferred protocol.

This protocol may be used to produce  $^{15}\text{N}$ -labeled biomolecules. Due to the fact that the only source of nitrogen is urea, this media lends itself to being a very cost-effective way to label proteins (the cell and all of its components as well) with  $^{15}\text{N}$ . The one caveat is that the host organism must be able to grow and produce the target protein in a defined media. A preferred host is *Bacillus subtilis*. Purification is made easier because the unwanted proteins are usually at level(s) lower than the target protein reducing the amount of contaminants to separate from this protein. The protocol is as follows:

#### 1) Media Preparation, Innoculation and Growth

These are the media and shake flask conditions preferred in the preparation of labeled biopolymers.

# **MOPS Medium-10X Bas for 1.0 L volum**

To a Milli-Q rinsed beaker add with stirring:

Milli-Q water	750mL
MOPS	83.72gm
Tricine	7.17gm
KOH Pellets	12.00gm
K <sub>2</sub> SO <sub>4</sub> (Potassium Sulfate) 0.276M Stock	10.00mL
MgCl <sub>2</sub> (Magnesium Chloride) 0.528M Stock	10.00mL
NaCl (Sodium Chloride)	29.22gm
Micronutrients - 100X Stock (previously made; recipe below)	100.00mL

5

Dissolve MOPS and Tricine, then add KOH. Add the remaining ingredients. Adjust the pH of the solution to 7.4 by addition of more KOH pellets (don't use a KOH solution as that could effect the final volume >1L). Generally ~2.13gm of additional KOH pellets are needed, be careful to ensure all KOH is solubilized before making additions of KOH pellets. With the pH at 7.4 adjust the liquid volume to 1.0L with additional Milli-Q water and after allowing the solution to mix well sterile-filter through a 0.22um filter unit.

10

Refrigeration of this media will help storage life, but it has been found that after ~1.5 to 2 months the MOPS media production level (for protease) decreases.

15

## **100X Micronutrients 1.00 liter**

Add the following ingredients, sequentially, to 1L Milli-Q water mix to solubilize then sterile filter through a 0.22µm filter unit. (Note: the actual volume will be 1.02L)

20

FeSO <sub>4</sub> *7H <sub>2</sub> O (Ferrous Sulfate, Heptahydrate)	400mg
MnSO <sub>4</sub> *H <sub>2</sub> O (Manganese Sulfate, Monohydrate)	100mg
ZnSO <sub>4</sub> *7H <sub>2</sub> O (Zinc Sulfate, Heptahydrate)	100mg
CuCl <sub>2</sub> *2H <sub>2</sub> O (Cupric Chloride, Dihydrate)	50mg
CoCl <sub>2</sub> *6H <sub>2</sub> O (Cobalt Chloride, Hexahydrate)	100mg
NaMoO <sub>4</sub> *2H <sub>2</sub> O (Sodium Molybdate, Dihydrate)	100mg
Na <sub>2</sub> B <sub>4</sub> O <sub>7</sub> *10H <sub>2</sub> O (Sodium Borate, Decahydrate)	100mg
CaCl <sub>2</sub> (Calcium Chloride) 1M Stock	10mL
C <sub>6</sub> H <sub>5</sub> Na <sub>3</sub> O <sub>7</sub> *2H <sub>2</sub> O (Sodium Citrate, Dihydrate) 0.5M Stock	10mL

**Shake Flask Media:** (For 1L volume)

10X Mops	100mL
21%Glucose/35% Maltrin M150 stock solution	100mL
<sup>15</sup> N-labeled Urea( <sup>15</sup> N <sub>2</sub> Urea,99 Atom%)	3.6gm
K <sub>2</sub> HPO <sub>4</sub> (Potassium Phosphate, DiBasic)	523mg
dH <sub>2</sub> O	

Mix the above ingredients and add deionized H<sub>2</sub>O to 1L volume. Mix well  
 5 and adjust the pH to 7.3(or predetermined best production pH between 7.0 to 7.5) with 50%NaOH. Add antibiotic(s) to desired concentration (e.g., 1mL of a 25mg/mL chloramphenicol (Cmp) solution added to this volume will give a 25ppm Cmp concentration) Sterile filter through a 0.22µm filter unit.

10 **Shake Flask conditions:** Using sterilized (e.g., autoclaved) shake flasks(bottom baffled are best for aeration of culture) use a 10 to 20% liquid volume(eg 50mL in a 250mL shake flask or 300mL in a 2800mL Fernbach)). For example, for protease production a 10 to 15% volume works well, for amylase production a 20% volume works well.

15 **Inoculation and Growth:** Cultures should be inoculated from thawed and mixed glycerol stocks (which were made in the Mops/Urea media prior to the labeling experiment) at the level of 150µL per 250mL shake flask or 1 vial(1.5mL) per 2800mL shake flask. Once inoculated the cultures should be grown at 37°C and 325  
 20 to 350rpm for ~60hrs (spo- host, cutinase production), ~72hrs (spo- host) for protease production and ~90hrs (spo+ host or amylase production), to achieve a maximum yield.

2) **Harvesting the culture(s)** Once the titers have reached their optimum level (or  
 25 reasonably close as predetermined in earlier experiments) the cultures should be harvested as the titers will only decrease and background biopolymers and by products will make the purification/isolation more difficult. Remove the shake flasks from the incubator and measure the activities from each culture (along with O.D. and pH). If all the activities are at a desirable level the cultures are pooled, and the pH is  
 30 adjusted to ~6.0 with acetic acid, (add slowly so that the resulting pH doesn't drift lower than the target pH). Centrifuge the broth immediately using centrifuge bottles appropriate for the amount of culture broth obtained. The material may be centrifuged at a high rpm (e.g., 12,000 rpm for 250mL bottles) for 30 minutes. Filter



the supernatants through 0.8 micron filters (Nalgene or Corning 1L units are preferred). Measure the total titer of this supernatant. The cell pellets can be saved, stored at -70°C, and used in future experiments as all of this material is labeled with  $^{15}\text{N}$ .

5

3) **Concentrating the Supernatant** This step should be done in a cold room (4°C) to minimize recovery loss. Use 400mL stirred cell(s) (Amicon 8400 series, 76mm diameter membranes) with a 10,000MWCO membrane (PM, polysulfone, is best, but may retain hydrophobic molecules). Add 350mL of the supernatant to each of the stirred cells, it is assumed that at least 1000mL of supernatant is available. Cap the units with their appropriate top and connect to a nitrogen line (50psi input), open the pressurizing valve on the unit and start concentrating. These units should be put on a multicell stir plate with ~130rpm stirring action. Add more supernatant to the cell(s) as the level goes down in the cell (usually 50-100mL at a time), make sure to collect the permeate in an appropriate beaker in case of a leak through the membrane. When all of the supernatant has been concentrated to at least one-tenth the original volume (e.g., 3000mL concentrated to 300mL) stop concentrating the material. Remove all the liquid from each stirred cell to a graduated cylinder, making sure to rinse the sides, stir bar and membrane off with a minimal amount of deionized water. This volume should be measured and an (activity) assay done to check the concentration of the labeled protein so that the total labeled protein available can be calculated (assays can be done on the permeate(s) to check for loss, also this material can be frozen away because all the protein components are labeled).

4) **Dialyzing the Concentrated  $^{15}\text{N}$  Biopolymer** If the first step in purifying the labeled protein will be ion-exchange the concentrated material should be dialyzed into an appropriate buffer system (if not the sample is ready to be run using the desired chromatographic method/system that will give the best yield of pure  $^{15}\text{N}$  biopolymer). This is set up with dialysis tubing of 10,000MWCO (SpectraPor 7, 32mm), filling the tubing with the concentrate, never more than 75mL per tube, clamping off the set up and put into a graduated cylinder (in the 4°C cold room) filled with buffer (20mM MES, pH 5.5, 1mM  $\text{CaCl}_2$  works well for most applications) on a stir plate (slowly stirring). The quantity of buffer used is between 20 to 50 times the volume of concentrate being dialyzed, and fresh buffer should be used after 4hours to ensure a good dialysis. It works best to let the sample dialyze overnight in the second buffer exchange. When done the sample should be removed from the dialysis tubing very carefully so that all the protein is recovered. At this point the



sample should be filtered with a 0.45micron filter unit, activity assays should be done along with a volume measurement.

5) **Purification of the  $^{15}\text{N}$  Biopolymers** As with any separation method one should know about the biopolymer that one is working with, because with this information it is easier to exploit specific characteristics of the molecule such as PI, hydrophobicity, affinity or any property that will distinguish it from the others in the media. For example, ion-exchange chromatography is the preferred method used to separate the labeled proteins from their matrix and works best if the PI of the target protein is known. Essentially the two pH ranges we have worked with so far is either pH 6.0 or pH 8.0, this involves using a cation exchange resin for binding the target protein and a salt (NaCl) gradient for elution of this protein. For good separation the load onto the column should be 25 to 35 per cent of the total column capacity, a 25cv (column volume) wash with the running buffer and a 50 to 100cv elution gradient where the eluate is collected in fractions. This ensures that the majority of the contaminants are eliminated from the protein sample fractions which will be pooled and assayed. At this point the pool is concentrated using a stirred cell in the cold room (4°C) and buffer exchanged/diafiltered to make another run using the either the same chromatographic procedure or a complimentary procedure involving conservative fractionation of the eluate. It is here that the pooled target biopolymer should be buffer exchanged while concentrating the sample in the buffer system that will be used for sample storage, whether frozen at minus20°C or formulated for future use. The amount of concentration of the sample is determined by the desired final biopolymer concentration that is needed in future use.

6) **Analysis of the  $^{15}\text{N}$ -Biopolymer Sample for Future Reference** Prior to the generation of the labeled biopolymer a pure sample of this unlabelled biopolymer should have been produced and well characterized by appropriate means. For example, for proteins SDS Page gel, activity assay, protein assay (e.g., BCA titration), amino acid analysis and a tryptic digest/peptide map along with MS analysis should have been done numerous times. With this information in hand the analysis of the labeled biopolymer is greatly facilitated as it is used for comparison to standardize the labeled biopolymer. All the analysis that was done for the unlabelled biopolymer should be done for the labeled biopolymer and compared the unlabelled biopolymer in different concentration ratios.

Purification and Calibration of Proteins and Peptides

The target biopolymer or analog, produced in isotope-labeled form either by synthesis or *in vivo*, can be purified by any means known in the art. For example, some extracellular alkaline proteases of microbial origin can be obtained in pure form by a single cation exchange chromatography step at pH 7.8 to 8.0 (Christianson and Paech, 1994). Other extracellular alkaline proteases can be obtained in pure form by cation exchange chromatography at pH 5.5 to 5.8 (Hsia et al., 1996), and yet other enzymes and proteins can be purified using one or more similar or different separation techniques, such as anion exchange, affinity, or hydrophobic interaction chromatography, size-exclusion chromatography, chromatofocusing, preparative isoelectrofocusing, precipitation, ultrafiltration, and others (for overviews see Deutscher, 1990, Scopes, 1994, and Janson and Rydén, 1998).

Peptides of specific sequence can be synthesized by standard techniques, purified by reverse-phase chromatography (RP-HPLC).

Once the protein or peptide is purified, a proof of purity can be ascertained, e.g. by SDS-PAGE for proteins, by RP-HPLC for peptides, the protein or peptide concentration can be determined by quantitative amino acid analysis, by total nitrogen analysis, by weight, or by light absorbance of the denatured protein (provided the amino acid sequence is known). Herein, a solution of purified protein or peptide of known protein mass content is called a 'calibrated solution'. The solution can be stabilized, as desired, by refrigeration, freezing, or by additives such as polyols and saccharides (1,2-propanediol, glycerol, sucrose, etc.), salt (sodium chloride, ammonium sulfate, etc.), and buffers adjusted to the pH of optimal stability.

Fragmentation of Proteins

The activity used in the practice of the present invention to fragment a protein into smaller fragments can be any enzyme or chemical activity which is capable of repeatedly and accurately cleaving at particular cleavage sites. Such activities are widely known and a suitable activity can be selected using conventional practices. Examples of such enzyme or chemical activities include the enzyme trypsin which hydrolyzes peptide bonds on the carboxyl side of lysine and arginine (with the exception of lysine or arginine followed by proline), the enzyme chymotrypsin which hydrolyzes peptide bonds preferably on the carboxyl side of aromatic residues (phenylalanine, tyrosine, and tryptophan), and cyanogen bromide (CNBr) which chemically cleaves proteins at methionine residues. Trypsin is often a preferred enzyme activity for cleaving proteins into smaller pieces, because trypsin is



characterized by low cost and highly reproducible and accurate cleavage sites. Techniques for carrying out enzymatic digestion are widely known in the art and are generally described by Allen, 1989, Matsudaira, 1993, Hancock, 1996, and Kellner et al., 1999.

5           Fragmentation of Polynucleotides

The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors and other requirements would be known to the ordinarily skilled artisan. For analytical purposes, typically 1 µg of plasmid or DNA fragment is used with about 2 units of enzyme in about 20 µl of buffer solution. For  
10 the purpose of isolating DNA fragments, typically 5 to 50 µg of DNA are digested with 20 to 250 units of enzyme in a larger volume. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturer. Incubation times of about 1 hour at 37° C are ordinarily used, but may vary in accordance with the supplier's instructions. After digestion the reaction is  
15 electrophoresed directly on a polyacrylamide gel to isolate the desired fragment.

Peptide Resolution

Any suitable separation technique can be used to resolve the peptide fragments. In one embodiment, a chromatographic column is employed comprising a chromatographic medium capable of fractionating the peptide digests as they are  
20 passed through the column. Preferred chromatographic techniques include, for example, reverse phase, anion or cation exchange chromatography, open-column chromatography, and high-pressure liquid chromatography (HPLC). Other separation techniques include capillary electrophoresis, and column chromatography that employs the combination of successive chromatographic techniques, such as  
25 ion exchange and reverse-phase chromatography. In a further embodiment, precipitation and ultrafiltration as initial clean-up steps can be part of the peptide separation protocol. Methods of selecting suitable separation techniques and means of carrying them out are known in the art. Herein, precipitation, ultrafiltration, and reverse-phase HPLC are preferred separation techniques.

30           Polynucleotide Resolution

Any suitable separation technique can be used to resolve the polynucleotide fragments. In one embodiment, size-based analysis of polynucleotide samples relies upon separation by gel electrophoresis (GEP). Capillary gel electrophoresis (CGE) may also be used to separate and analyze mixtures of polynucleotide fragments  
35 having different lengths, e.g., the different lengths resulting from restriction enzyme cleavage. In a preferred embodiment, the polynucleotide fragments which differ in

11/01/2006 10:00 AM GC626-2app 20



base sequence, but have the same base pair length, are resolved by techniques known in the art. For example, gel-based analytical methods, such as denaturing gradient gel electrophoresis (DGGE) and denaturing gradient gel capillary electrophoresis (DGGC), can detect mutations in polynucleotides under "partially denaturing" conditions. Recently, a Matched Ion Polynucleotide Chromatography (MIPC) separation method has been described for the separation of polynucleotides. See U.S. Patent No. 6,265,168.

#### Mass Spectrometric Identification of Peptides

Any suitable mass spectrometry instrumentation can be used in practicing the present invention, for example, an electrospray ionization (ESI) single or triple-quadrupole, or Fourier-transform ion cyclotron resonance mass spectrometer, a MALDI time-of-flight mass spectrometer, a quadrupole ion trap mass spectrometer, or any mass spectrometer with any combination of source and detector. A single quadrupole and an ion-trap ESI mass spectrometer are especially preferred herein.

#### General Embodiments/Examples

As used herein, "percent homology" of two amino acid sequences or of two nucleic acid sequences is determined using the algorithm of Karlin and Altschul (Proc. Natl. Acad. Sci. USA 87:2264-2268, 1990), modified as in Karlin and Altschul (Proc. Natl. Acad. Sci. USA 90:5873-5877, 1993). Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul et al. (J. Mol. Biol. 215:403-410, 1990). BLAST nucleotide searches are performed with the NBLAST program, score = 100, wordlength = 12, to obtain nucleotide sequences homologous to a nucleic acid molecule of the invention. BLAST protein searches are performed with the XBLAST program, score = 50, wordlength = 3, to obtain amino acid sequences homologous to a reference polypeptide. To obtain gapped alignments for comparison purposes, Gapped BLAST is utilized as described in Altschul et al. (Nucleic Acids Res. 25:3389-3402, 1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) are used. See <http://www.ncbi.nlm.nih.gov>.

A biopolymer or biopolymer fragment is said to "correspond" to an analog thereof when the biopolymer/fragment and analog have similar chemical and physical properties, but differ in at least one chemical or physical property. For example, an analog of a target polypeptide can comprise a polypeptide having an amino acid sequence identical to that of the target, the analog being formed, however, from amino acids that differ isotopically from those making up the target polypeptide. Or, the polypeptide analog can be isotopically identical to the target in

terms of its amino acid content, but have an amino acid sequence that is homologous, but not identical, to the sequence of the target (e.g., the analog can have one or more amino acid substitutions, insertions, or deletions (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 substitutions)). In one embodiment, the analog shares at least 90,  
5 95, and/or 98 percent homology with the target biopolymer. Alternatively, the analog can be derivatized (e.g., tagged) in a fashion so as to alter at least one chemical or physical property as compared to the target. The exact manner in which the analog differs from the biopolymer is not critical, provided only that the two are capable of producing a pair of peaks that can be distinguished one from the other, yet which  
10 occur relatively close to one another, in mass spectrographic analysis (i.e., a peak pair can be identified attributable to the target and analog).

#### Known Protein

In one embodiment of the present invention, which is especially useful for the analysis of a known protein or a family of proteins that share a high degree of  
15 sequence homology with the known protein as in the case of genetically modified variants of a parent molecule, or closely related molecules with the same function, but from different organisms, (e.g., having at least 85%, 90%, 95%, and/or 98% sequence homology) a purified, isotope-labeled, calibrated form (analog) of a target protein is added to a solution (e.g., a cell extract) known or believed to contain the  
20 target protein. The resulting mixture is subjected in its entirety to rapid protein fragmentation, e.g., by trypsin digestion. The resulting peptides are briefly separated, e.g., by reverse-phase chromatography, and the eluting peptides are monitored by mass spectrometry. The ratio of integrated peak areas of a reconstructed ion current chromatogram of corresponding peptides (wildtype and  
25 isotope-labeled) provides a direct measure for the molar concentration of the unknown concentration of the known protein.

As detailed in Example 1, the inventors have tested such a method with <sup>15</sup>N-*Bacillus lentus* subtilisin-N76D-S103A-V104I (<sup>15</sup>N-subtilisin-DAI), and accurately  
30 determined the unknown concentrations of subtilisin-DAI to ±5%. In other experiments, correct concentrations were obtained with a standard-to-target mass ratio of up to 10:1, with as low as 2 µg · ml<sup>-1</sup> and as little as 2 µg of target protein (see Table II). In yet another experiment, the fragmentation time was reduced to 1 min, and the total chromatography cycle was limited to 20 min (see Figure 3).

The technique has been validated by using the same internal standard for a  
35 large number of variants with as many as ten different mutations, some of which affect the catalytic properties so that rate measurements could not serve as a



convenient or reliable way of quantifying the proteins in crude solutions. With an extended chromatography regime, one can pinpoint the approximate area of mutation, and in some cases even the exact mutation. It should be appreciated that there is no limit to the sequence variation as long as at least one peptide is shared  
5 between the internal standard and the target protein. The application of the methods of the present invention to the quantitation of variants that have lost catalytic function is of particular interest. In one specific case, this technique was used to quantitate a putative alkaline serine protease in a commercially available, solid fermentation product, as detailed in Example 2.

#### 10 Unknown Protein

The methods of the present invention can be applied to unknown (putative) polypeptides, as well. Analysis of such polypeptides can be accomplished, for example, using synthetic isotope-labeled peptides, or by calibrating an isotope-labeled cell extract with peptides of natural abundance atomic composition. In an  
15 embodiment of the latter, a putative protein of interest is selected using one or more available databases and software tools. A number of sequence libraries can be used, including, for example, the GenBank database (now centered at the National Center for Biotechnology Information, Bethesda, summarized by Burks et al., 1990), EMBL data library (now relocated to the European Bioinformatics Institute,  
20 Cambridge, UK, summarized by Kahn and Cameron, 1990), the Protein Sequence Database and PIR-International (summarized by George et al., 1996), and SWISS-PROT (described in Bairoch and Apweiler, 2000). The ExPASy (Expert Protein Analysis System) proteomics server of the Swiss Institute of Bioinformatics (SIB), at <http://www.expasy.ch/>, provides information on, and URLs (links) for, numerous  
25 available databases and software tools for the analysis of protein sequences. Another listing of URLs to access tools for protein identification and databases on the Internet is set out by Lahm and Langen, 2000.

For example, in a case where it is desired to select a putative protein of a Bacillus species, one can search a database of Bacillus sequence information, e.g.,  
30 as described by Kunst et al., 1997, and available over the Internet at <http://genolist.pasteur.fr/SubtilList/>. It should be appreciated that the present invention is applicable to any sequence databases and analysis tools available to the skilled artisan, and is not limited to the examples described herein.

Once a putative protein has been selected, a theoretical fragmentation (e.g.  
35 trypsin digest) of the protein of interest is performed. Several programs to assist with protease digestion analysis are available over the Internet. MS-Digest, for



example, (available at <http://prospector.ucsf.edu/>) allows for the "in silico" digestion of a protein sequence with a variety of proteolytic agents including trypsin, chymotrypsin, V8 protease, Lys-C, Arg-C, Asp-N, and CNBr. The program calculates the expected mass of fragments from these virtual digestions and allows  
5 the effects of protein modifications such as N-terminal acetylation, oxidation, and phosphorylation to be considered. From the theoretical fragmentation, a suitable peptide is selected, which can then be synthesized and calibrated. The suitability of the peptide can be checked by querying the genome of interest for redundancy. If the same peptide (string of amino acid residues) occurs on more than one protein  
10 then another peptide should be selected.

Next, the organism can be grown on isotope-enriched media. In a preferred embodiment, the nitrogen content of the media is enriched in  $^{15}\text{N}$ . The calibrated peptide is added to a protein extract from the cells, and the entire mixture is digested rapidly and 'cleaned up'; for example, and without limitation, by precipitation, ultra-  
15 filtration, or ion exchange chromatography. The choice of an optimal technique can be tailored by the skilled artisan to the properties of the peptide (size, charge, hydrophobic index, etc.) since these features can be established prior to the use of the peptide as an internal standard. The resulting 'lean' solution is passed over a RP-HPLC column attached to a mass spectrometer. Since the characteristics of the  
20 internal standard peptide (retention time, mass) are known, the skilled artisan can focus the separation and the mass measurement on a very narrow window, both in time and mass, and thereby tremendously increase the sensitivity of the detection. If the expected peak pair is found (wild-type from internal standard,  $^{15}\text{N}$  from organism), peak area integration yields the absolute concentration of the targeted  
25 protein. Preferably, in this embodiment, a series of experiments is carried out, as appropriate, to assure that the fragmentation of the target protein is substantially complete with respect to the peptide of interest. The  $^{15}\text{N}$ -labeled extract can be queried for any number of proteins, even simultaneously, as long as mass and retention times can be properly spaced.

Advantageously, the just-described method provides a calibrated  $^{15}\text{N}$ -labeled  
30 protein mixture (cell extract) that can be conserved (e.g., in small aliquots) for later use. For example, now possessing a calibrated  $^{15}\text{N}$ -labeled cell extract, the organism can be grown under defined conditions, and extracts queried for the presence, for an increase or decrease of the absolute concentration of the target  
35 protein by mixing it with the calibrated  $^{15}\text{N}$ -labeled aliquot. It should be appreciated that, at this stage, the digest does not have to be quantitative as long as a little of the





fragment of the molecule of interest is formed. Analysis can be carried out by LC/MS as above. The skilled artisan can increase the accuracy of absolute quantitation by searching for one or more other peptides from the target protein because they all must exist as pairs. A byproduct of this approach is that any protein other than the target proteins can be quantified relative to the level in the isotope-labeled sample similar to the approach taken by others using isotope labeling (Oda et al., 1999) and reporter groups (Gygi et al., 1999).

Additional General Embodiments/Examples

The teachings herein can be adapted to a number purposes. For example, the selected target can be a polymer of nucleotides, e.g., one or more polynucleotides and/or oligonucleotides. According to one general embodiment, a target oligonucleotide is selected for analysis and an analog of the target oligonucleotide is generated. The target oligonucleotide can be, for example, an oligonucleotide that is known to be in a mixture, a putative oligonucleotide (e.g., derived from a genome database search) that is potentially present in a mixture, or a known or putative oligonucleotide segment or fragment. The analog of the target oligonucleotide can be the target oligonucleotide itself or a unique segment or fragment of the target oligonucleotide. One or the other of the target oligonucleotide and analog is labeled, using methods known in the art (e.g., <sup>32</sup>P labeling), so that the two can be distinguished from one another in subsequent mass analysis. The analog is purified and its absolute quantity is determined in a solid quantity or in a solution by standard techniques (the analog is now said to be 'calibrated'), and a known amount is employed as an internal standard in the solution to be assayed. The oligonucleotides of the mixture are treated with a fragmenting activity (e.g., an endonuclease), and the oligonucleotide fragments of the mixture are then resolved. Corresponding oligonucleotide fragment pairs are then analyzed by selected ion monitoring of a mass spectrometer. Peak area integration of such pairs provides a direct measure for the amount of target oligonucleotide in the crude solution.

The present teachings can be adapted for the identification of a target biopolymer fragment in a crude solution or mixture. In one embodiment, wherein a fragment of a target protein is identified in a solution otherwise not including such fragment (i.e., the fragment to be identified is not natively present in the solution), a selected fixed ratio of an analog of the target protein and the target protein are added to the solution. The target protein and analog are then subjected to fragmentation, e.g., by treatment with a fragmenting activity, thereby generating a plurality of corresponding peptide pairs. The peptide fragments are then resolved,



e.g., by way of a suitable chromatographic technique. Mass spectrometric analysis is then employed to identify those fragment pairs corresponding to the target protein that exhibit the selected ratio. In other words, the fragments that arose from the target protein are identified via their characteristic (selected) mass ratio. Next, the fragment pairs exhibiting the selected ratio can then be sequenced using any suitable technique, e.g., utilizing further mass spectrometric analysis, database query, etc. (see, e.g., Lahm and Langen, 2000; Corthals et al., 1999).

The following preparations and examples are given to enable those skilled in the art to more clearly understand and practice the present invention. They should not be considered as limiting the scope and/or spirit of the invention, but merely as being illustrative and representative thereof.

In the experimental disclosure which follows, the following abbreviations apply: eq (equivalents); M (Molar);  $\mu$ M (micromolar); N (Normal); mol (moles); mmol (millimoles);  $\mu$ mol (micromoles); nmol (nanomoles); g (grams); mg (milligrams); kg (kilograms);  $\mu$ g (micrograms); L (liters); ml (milliliters);  $\mu$ l (microliters); cm (centimeters); mm (millimeters);  $\mu$ m (micrometers); nm (nanometers); ° C. (degrees Centigrade); h (hours); min (minutes); sec (seconds); msec (milliseconds); Ci (Curies) mCi (milliCuries);  $\mu$ Ci (microCuries); TLC (thin layer chromatography).



## EXAMPLES

The following examples are illustrative and are not intended to limit the invention.

### Example 1

#### 5 1A. Materials and Methods

*Bacillus lentus* subtilisin-N76D-S103A-V104I (subtilisin DAI) was expressed by *Bacillus subtilis* grown on minimal media and  $^{15}\text{N}$ -urea as nitrogen source. The protein was purified (Goddette et al., 1992; Christianson and Paech, 1994) and calibrated by amino acid analysis and by active site titration (Hsia et al., 1996) as described previously. Once calibrated, succinyl-L-alanyl-L-alanyl-L-prolyl-L-phenylalanyl-*p*-nitroanilide (sucAAPF-pNA) supported catalytic activity in 0.1 M Tris/HCl, containing 0.005% (v/v) Tween 80, pH 8.6 at 25°C, recorded at 410 nm and measured in  $\text{AU} \cdot \text{min}^{-1}$ , was used to quantify the enzyme concentration ( $f = 0.020 \text{ mg} \cdot \text{min} \cdot \text{AU}^{-1}$ ). Wildtype *Bacillus lentus* subtilisin (subtilisin) was purified, 15 calibrated, and measured similarly ( $f = 0.053 \text{ mg} \cdot \text{min} \cdot \text{AU}^{-1}$ ).

Standard peptide mapping with trypsin was carried out as outlined by Christianson and Paech, 1994, except that sample sizes ranged from 2 to 100  $\mu\text{g}$  of protein. Peptides were separated by HPLC (Hewlett-Packard model 1090) on a  $\text{C}_{18}$  reverse-phase column (Vydac, 2.1x150 mm), heated to 50°C, using a gradient of 20 0.08% (v/v) trifluoroacetic acid (TFA) in acetonitrile and 0.1% (v/v) TFA in water. The column eluate was monitored by UV absorbance at 215 nm and by mass measurement on an ESI mass spectrometer (Hewlett-Packard, model 5989B/59987B).

Rapid peptide mapping was performed with a trypsin-to-protein ratio of 1:1 for 25 15 s to 1 min at 37°C. Peptides were separated on 2.0x50 mm  $\text{C}_{18}$  reverse-phase column (Jupiter, by Phenomenex).

#### 1B. Results

Figure 1: UV traces of a tryptic co-digest of  $^{15}\text{N}$ -subtilisin DAI and subtilisin, . 30 Peptides are numerated in the order of occurrence beginning with the N-terminus (see Table I).

Figure 2. (A) Integrated total ion current (TIC) chromatogram of peptide 3 of subtilisin (indexed (s)) and  $^{15}\text{N}$ -subtilisin DAI (indexed ( $^{15}\text{N}$ )). (B) TIC of peptides 5, 6 35 and 9 of  $^{15}\text{N}$ -subtilisin DAI and subtilisin. The results of area integration for both TIC and UV peaks are summarized in Table I. Note that sequence differences of

subtilisin and subtilisin-DAI reside on peptide 5 (N74D) and 6 (S101I, V102A).  
Amino acid sequence numbering is linear.

Table I.: Sequence comparison, m/z values, and ratios of integrated TIC peak areas  
and UV absorbance peak areas for chromatograms in Figure 1. The concentration  
measured by the co-digest technique for subtilisin and subtilisin-DAI was 8.15 and  
7.13 mg/ml, respectively, while the given concentration (established by independent  
methods) was 7.99 and 7.03mg/ml, respectively.

#### Example 2

A fermentation broth concentrate of unknown origin was suspected of  
containing an alkaline serine protease. A small sample was dissolved in buffer and  
spiked with purified <sup>15</sup>N-labeled subtilisin-Y217L. The mixture was digested with  
trypsin, peptides were separated by RP-HPLC, and the eluate monitored by UV  
absorbance and by mass spectrometry. Figure 4 (A) shows an SDS-PAGE gel of  
the composition of the sample. Figure 4 (B) displays the peptide map, and Figure 5  
gives a few examples of TIC traces. The data show that the sample contains an  
alkaline serine protease closely related to subtilisin BPN', and in this case,  
specifically at 0.54 mg· ml<sup>-1</sup>.

### Example 3

Randomly generated variants of subtilisin-DAI were expressed by cultures grown on minimal media in microtiter plates. Aliquots of cell-free supernatants were probed for the presence of subtilisin-DAI variants by co-digests with  $^{15}\text{N}$ -labeled subtilisin-DAI. In separate experiments the catalytic activity was measured. In yet another experiment, the ratio of specific concentration to activity (referred to as 'conversion factor' f) was measured by active site titration with a mung bean inhibitor (MBI) solution calibrated in the same experiment with a previously standardized solution of subtilisin-DAI (Hsia et al., 1996). The data shown in Table II show convincingly the accuracy of the peptide mapping method for protein concentration measurements. A further advantage of the technique is that the protein variants can be queried for similarities and approximate location of mutations. Because all peptides of the internal standard are known, each can be checked for the presence of the unlabeled counterpart. If not present the target protein has a mutation on that sequence. Next one would search for a peptide of closely related mass and verify that it exists in the quantity, anticipated from the quantity of those peptides identical in sequence with the internal standard, using the UV trace.

### Example 4

From the previous example one can extrapolate that the method should work with equal efficiency and accuracy for proteins of unknown properties but known sequence by using instead of purified  $^{15}\text{N}$ -labeled protein a synthetic  $^{15}\text{N}$ -labeled peptide. This will be added to the sample ready for trypsin digestion. After digestion the sample will be analyzed as before.

### Example 5 $^{15}\text{N}$ Protease

This example describes a method for the batch preparation of a  $^{15}\text{N}$ -labeled protease. The Mops/Urea shake flask protocol (described above) was used with all of the chemicals, except for the urea, purchased from Sigma chemical in highest purity available.  $^{15}\text{N}_2$  Urea(99 atom%) was purchased from Isotec, Inc. A 1.8L batch of media was prepared with chloramphenicol at 25ppm and sterile filtered. 300mL was added aseptically to each of the 6 sterilized 2.8L bottom baffled fernbachs. The inoculation was done by adding the thawed and mixed glycerol stocks, protease hyper producer prepared previously in the Mops/urea media and frozen, at 1vial(1.5mL) per shake flask. The shake flasks were put into a New Brunswick shaker/incubator, after inoculation, and run at 37°C and 350rpm for 78hours. At the harvest point, 78hours, AAPF activity assays were done on the samples and titers

ranged from 0.7g/L to 1.4g/L. The contents from the shake flasks were pooled together, pH adjusted to 5.5 with acetic acid and centrifuged in 250mL bottles at 12,000rpm for 30minutes. The supernatants were filtered with a 0.8 micron Nalgene 1L filter unit. The pool was assayed at 1.1g/L for 1700mL with the total <sup>15</sup>N protease being 1.9gms. The supernatant was concentrated in the cold room (@4°C) to 135mL, using 3 Amicon 8400 stirred cells and PM10 (10,000MWCO) membranes. There was no loss of protein in the concentration step.

Dialysis was done using 20mM MES, pH 5.4, 1mM CaCl<sub>2</sub> buffer in a 15L graduated cylinder on a stir plate in the cold room, with the sample being added in two 67.5mL aliquots respectively to 10,000MWCO Spectra Por 7 dialysis tubing, clamped off and placed into the cylinder with buffer. After the overnight dialysis the samples were removed from the graduated cylinder, the clamps removed from the dialysis tubing and the contents poured into and filtered using a 0.45micron Nalgene 500mL filter unit. Assays run at this time showed no loss of protein at 1.9gm total available in 250mL.

The protease protein was purified using a low pH buffer system with a cation exchange column because the PI of the enzyme is around 8.6. An Applied Biosystems Vision was used to do the purification along with a 16x150mm (32mL) column of POROS HS 20 (Applied Biosystems cation exchange resin). The program used to do the purification is as follows: Equilibrate the column at 50mL/minute with 20cv's (colume volumes) of 20mM MES, pH 5.4, 1mM CaCl<sub>2</sub> buffer, load the sample (150mL) onto the column at 15mL/minute, wash the column at 50mL/minute with a gradient from the 20mM MES, pH 5.4, 1mM CaCl<sub>2</sub> buffer to 20mM MES, pH 6.2, 1mM CaCl<sub>2</sub> buffer in 25cv's. Elute the <sup>15</sup>N protease protein with a gradient from 20mM MES, pH 6.2, 1mM CaCl<sub>2</sub> buffer to 20mM MES, pH 6.2, 1mM CaCl<sub>2</sub>, 15mM NaCl buffer in 75cv's(start collecting the fractions at 5cv's into the gradient). Finally, clean the column off with a salt wash of 2M NaCl 10cv's, rinse with 10cv's of H<sub>2</sub>O. This run was made three times to purify all of the labeled protein, the <sup>15</sup>N protease came off the column between 8 to 12mM NaCl, with 95 11mL fractions collected each run. The labeled protease was concentrated from 1.8L to 150mL using an Amicon stirred cell with a 10,000MWCO PM membrane, with a buffer exchange/diafiltration to 20mM MES, pH 5.4, 1mM CaCl<sub>2</sub> to prepare the sample for another run on the same system with the same method. Some of the labeled protease was lost because of the cuts made on the fractions collected, with the total available <sup>15</sup>N protease down to 1.4gm. After three more runs the purification was done. There was a pool of purified material with a 1.3L total volume. This was

concentrated down to 65mL using the Amicon concentrator and a buffer exchange to 20mM MES, pH 5.4, 1mM CaCl<sub>2</sub> buffer. The <sup>15</sup>N protease purified sample was sterile filtered through a 0.22micron using the Nalgene 0.22micron 250mL filter unit. An AAPF activity assay showed the concentration to be 20g/L (mg/mL) and this was aliquoted into 60 Nalgene 1.8mL cryovials at 1mL of sample each (the identity, date and concentration was labeled onto each vial). These vials were frozen at -20°C in a labeled container.

Analysis was done on these samples to confirm the concentration, the purity and the presence of the <sup>15</sup>N labeling. An SDS-PAGE gel run against an unlabelled protease standard showed no molecular weight bands greater than 27,480, the intensity of the protease bands at 27,480 Daltons was about the same with the subsequent breakdown bands (3) to be of the same intensity also. An amino acid analysis showed that the AAPF activity concentration to be the same (20g/L) as well as the BCA total protein concentration run against the unlabelled protease standard. Tryptic digests/codigests with protease (unlabelled) and subsequent peptide mapping with MS analysis on the HP 59987A engine showed that the peptides were labeled with <sup>15</sup>N. Thus, the material was shown to be what was intended, <sup>15</sup>N labeled protease, suitable for analytical use.

Those skilled in the art will appreciate the numerous advantages offered by the present invention. For example, unlike the prior methods, the methods taught herein can yield absolute protein concentrations. In comparison, ICAT (Gygi et al., 1999) measures relative quantities, as does staining of 2D gels or the isotope technique by Oda et al., 1999. A further advantage of the present method is that it applies to all proteins, while the ICAT technology can capture only about 10% of all proteins since it relies on the presence of free SH groups. Yet a further advantage of the present invention is that this methodology is compatible with all automated equipment developed for protein identification under the 'proteomics' umbrella.

The present invention is useful where only very dilute concentrations of biopolymer are available for analysis. With regard to quantity, for example, the present invention can be employed to determine the absolute quantity of a selected protein in a solution containing less than 25, less than 20, less than 15, less than 10, less than 5, and down to about 2 micrograms, or less, of such protein. With regard to concentration, the present invention can be employed to determine the absolute quantity of a selected protein in a solution containing less than 25, less than 20, less



than 15, less than 10, less than 5, and down to about 2 micrograms/ml, or less, of such protein.

Various other examples and modifications of the foregoing description and examples will be apparent to a person skilled in the art after reading the disclosure  
5 without departing from the spirit and scope of the invention, and it is intended that all such examples or modifications be included within the scope of the appended claims. All publications and patents referenced herein are hereby incorporated by reference in their entirety.

10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
220